

Viconin

Cefquinome sulfate Intramammary Infusion (Lactating Cow)

INDICATIONS

For the treatment of clinical mastitis in lactating cows caused by *Staphylococcus aureus*, *Streptococcus uberis*, *Streptococcus dysgalactiae*, *Escherichia coli* and other enterobacteria susceptible to cefquinome.

DOSAGE & ADMINISTRATION

The content of one syringe should be infused gently into the teat of the infected quarter every 12 hours after each of three successive milkings.

Milk out the affected quarter(s). After thoroughly cleaning and disinfecting the teat and teat orifice with the cleaning towel provided, gently infuse the contents of one syringe into each affected quarter. Disperse the product by gentle massage of the teat and udder of the affected animal.

CAUTIONS

*Handle this product with great care to avoid exposure, taking all recommended precautions.

*If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the Doctor this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention. Wash hands after using the cleaning towels and wear protective gloves if skin irritation due to Isopropyl alcohol is known or suspected.

*The syringe must only be used once. Partly used syringes should be discarded.

WITHDRAWAL TIME

Meat and offal, 2 days; Milk, 96 hours

STORAGE CONDITIONS

Store in an airtight container below 25 °C, protected from light.

PACKAGING SIZE

8.0 g/syringe.

Each 8.0 g (1 syringe) contains Cefquinome (as cefquinome sulfate) 75 mg.

