

Journal Pre-proof



GhUBX controlling helical growth results in production of stronger cotton fiber

Yihao Zang, Yan Hu, Chenyu Xu, Shenjie Wu, Yangkun Wang, Zhiyuan Ning, Zegang Han, Zhanfeng Si, Weijuan Shen, Yayao Zhang, Lei Fang, TianZhen Zhang

PII: S2589-0042(21)00898-1

DOI: <https://doi.org/10.1016/j.isci.2021.102930>

Reference: ISCI 102930

To appear in: *ISCIENCE*

Received Date: 17 January 2021

Revised Date: 9 July 2021

Accepted Date: 27 July 2021

Please cite this article as: Zang, Y., Hu, Y., Xu, C., Wu, S., Wang, Y., Ning, Z., Han, Z., Si, Z., Shen, W., Zhang, Y., Fang, L., Zhang, T., *GhUBX* controlling helical growth results in production of stronger cotton fiber, *ISCIENCE* (2021), doi: <https://doi.org/10.1016/j.isci.2021.102930>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 The Author(s).

1 ***GhUBX* controlling helical growth results in production of**
2 **stronger cotton fiber**

3 Yihao Zang^{1,2}, Yan Hu², Chenyu Xu^{1,2}, Shenjie Wu³, Yangkun Wang¹,
4 Zhiyuan Ning¹, Zegang Han¹, Zhanfeng Si², Weijuan Shen¹, Yayao Zhang²,
5 Lei Fang², TianZhen Zhang^{1,2,4,*}

6

7 ¹ State Key Laboratory of Crop Genetics and Germplasm Enhancement,
8 College of Agriculture, Nanjing Agricultural University, Nanjing 210095,
9 China.

10 ² Agronomy Department, College of Agriculture and Biotechnology,
11 Zhejiang University, Hangzhou 310029, China

12 ³ Biotechnology Research Center, Shanxi Academy of Agricultural
13 Sciences, Taiyuan 030031, China

14 ⁴ Lead contact

15 *Correspondence and requests for materials should be addressed to TZ
16 Zhang (cotton@njau.edu.cn)

17

18 **Summary**

19 Cotton fiber is an excellent model for studying plant cell elongation and cell wall
20 biogenesis as well, because they are highly polarized and use conserved polarized
21 diffuse growth mechanism. Fiber strength is an important trait among cotton fiber
22 qualities due to ongoing changes in spinning technology. However, the molecular
23 mechanism of fiber strength forming is obscure. Through map-based cloning, we
24 identified the fiber strength gene *GhUBX*. Increasing its expression, the fiber strength
25 of the transgenic cotton was significantly enhanced compared to the receptor W0 and
26 the helices number of the transgenic fiber was remarkably increased. Additionally, we
27 proved that GhUBX regulates the fiber helical growth by degrading the GhSPL1 *via*
28 the ubiquitin 26S-proteasome pathway. Taken together, we revealed the internal
29 relationship between fiber helices and fiber stronger. It will be useful for improving the
30 fiber quality in cotton breeding and illustrating the molecular mechanism for plant
31 twisted growth.

32 **Keywords:** QTL mapping, GhUBX, GhSPL1, helical growth, fiber strength, ubiquitin
33 26S-proteasome pathway

34

35

36 **Introduction**

37 Helical growth is a widespread phenomenon in the plant kingdom. It is seen in the
38 coiling of tendrils, reversal growth of buds and leaves, spiraled arrangement of petals,
39 and twisting of leaf blades. There are many examples of helical growth in plants, not
40 only in the tendrils of climbing plants such as the grapevine *Vitis vinifera* and the
41 pedicels of orchid buds (e.g., the *Cattleya* hybrid, family Orchidaceae), but also in the
42 petals of *Lagunaria patersonia* (family Malvaceae) and leaves of the geebung shrub
43 *Persoonia helix* (family Proteaceae) (Jaffe and Galston, 1968; Smyth, 2016). Generally,
44 the shape and movement of plants are determined by the directional expansion of cells,
45 which is caused by the interaction between the cell turgor pressure and cell wall tension
46 (Skotheim and Mahadevan, 2005; Dumais and Forterre, 2012). The spatial distribution
47 of cell wall microfibrils determines the polarity of the cell expansion shaping the plant
48 form. Multiple recent studies have highlighted the fact that the cortical microtubules
49 play an important role in microfibril orientation. For instance, membrane-assisted
50 cortical microtubules regulate the arrangement of cellulose microfibrils (Tiwari and
51 Wilkins, 1995; Baskin, 2005; Paredez et al., 2006). The microtubule and actin
52 cytoskeletons cooperate to influence shape change in plant cells (Yanagisawa et al.,
53 2015). In most situations, the helical growth of plant cells is associated with
54 rearrangement of cortical microtubules. These helical rearrangements have been
55 proposed to drive the handedness of cell elongation (Ishida et al., 2007).

56 Allotetraploid upland cotton (*Gossypium hirsutum* L.) is the leading natural fiber-
57 producing species, accounting for 95% of cotton production worldwide. Cotton fiber
58 cells show twisted growth as they grow. Li *et al.* (2009) reported that the overexpression
59 of fibroin, a silkworm gene, could increase fiber helices. Fiber strength (FS) refers to
60 the ability of a fiber to withstand a load before breaking. It not only depends on the
61 amount of cellulose contained in cell walls (a higher percentage of strength is attributed
62 to greater rigidity of cellulosic chains), but also on the frequency and distribution of
63 reversal and convolutional structures in the microfibril helix (Hsieh, 1999; Haigler,
64 2010). In recent years, the improvement of fiber quality has been driven primarily by

65 the advent of high-speed rotor spinning. The convolutional structure could increase the
66 binding force between the fibers and improve the yarn strength during high-speed rotor
67 spinning.

68 Until now, several hundred quantitative trait loci (QTLs) related to fiber quality traits
69 (e.g., fiber strength, length, uniformity, micronaire) have been mapped using various of
70 genetic segregated populations (Zhang et al., 2003; Mei et al., 2004; Rong et al., 2007;
71 Lacape et al., 2010). However, pinpointing the exact location of genes with a large
72 effect underlying the QTL is still a mammoth task. To our knowledge, map-based
73 cloning of the exact gene conferring the QTL in cotton has not been reported. In this
74 study, we cloned a causal gene, *GhUBX*, responsible for higher fiber strength using
75 map-based cloning of a major-effect QTL, *qFS-D3-1*. The GhUBX reduces the content
76 of the GhSPL1 protein in fiber cells *via* the ubiquitin 26S-proteasome pathway; it acts
77 as the bridge between ubiquitin and the plant-specific microtubule-associated protein
78 GhSPL1. Increased *GhUBX* transcripts could significantly increase the fiber helices and,
79 consequently, improve the fiber strength of transgenic cotton.

80

81 **Results**

82 **Map-based cloning of the fiber strength QTL *qFS-D3-1***

83 The major-effect QTL for fiber strength, named *qFS-D3-1*, originated from Prema and
84 was mapped on chromosome D03 (Chr. D03) (**Figure 1A**). It could be identified
85 simultaneously in four environments with a phenotypic variation ranging from 2.11%
86 to 14.71% with substantially high logarithm-of-the-odds (LOD) scores that fluctuated
87 from 5.46 to 14.58 (Ning et al., 2014). To fine-map and clone this QTL, four
88 recombinant inbred lines (RILs), RIL43, RIL98, RIL120 and RIL168, that derived from
89 (Prema × 86-1) RILs and contained the common *qFS-D03-1* locus, were backcrossed
90 with 86-1 respectively to produce four secondary mapping populations (**Figures S1A,**
91 **S1B and Table S2**). The *qFS-D3-1* was detected in these four F₂ populations and
92 anchored within a 23.5-centimorgan interval with six pairs of single-nucleotide
93 polymorphisms (SNPs) and three pairs of insertion-deletion (InDel) markers (**Figure**

94 **S1C)**. Using the newly developed simple sequence repeats (SSRs), SNPs, and InDel
95 markers in this interval (Wang et al., 2015), we delimited the *qFS-D3-1* within a 1.14-
96 cM interval between markers K5219 and K5221 with a LOD of 15.54 using the
97 segregating (four RILs × 86-1) F₂ comprised of 1,864 individuals (**Table S3**),
98 corresponding to a 0.93-megabase (Mb) physical distance on Chr. D03. Based on our
99 updated genome sequence of *G. hirsutum* acc. TM-1 (Hu et al., 2019), 23 genes had
100 been annotated in this short region (0.93 Mb) (**Table S4**). Transcriptomic data
101 (<https://cotton.zju.edu.cn>) showed that 11 of 23 candidate genes were expressed during
102 the stages of fiber development and confirmed by the real-time quantitative polymerase
103 chain reaction (qRT-PCR) (**Figures S1D and S2A**). Among these 11 genes, five showed
104 differential expression in just one fiber development stage between Prema and 86-1,
105 and one gene (*GH_D03G0985*) always showed a lower expression in the fibers of
106 Prema than those of 86-1 (**Figure S2A**). The full-length coding regions of the six
107 differentially expressed genes from Prema and 86-1 were isolated and sequenced.
108 Sequence alignment indicated that five genes differentially expressed in just one fiber
109 development stage had the same protein sequence between Prema and 86-1. However,
110 we detected a 6-bp (GCCTCC) SSR variation conferring the deletion of two amino
111 acids (Ala-Ser) at the N-terminal of *GH_D03G0985* (**Figure 1B**) in Prema compared
112 with 86-1. Therefore, *GH_D03G0985*, annotated as a UBX (*GhUBX*), was the
113 candidate gene for *qFS-D3-1*.

114

115 **The SSR in *GhUBX* has a significant correlation with the fiber strength trait**

116 Sequence alignment showed that the haplotype of *GhUBX* gene was classified into the
117 *GhUBX*-Prema type, with CTCGGCCTCT (GCCTCC)₅, and the *GhUBX*-86-1 type,
118 with CTCGGCCTCT (GCCTCC)₆. The SSR marker K222 was designed surrounding
119 this SSR variation. The 183 RILs derived from (Prema × 86-1) were divided into two
120 major categories by the marker K222, *GhUBX*-Prema and *GhUBX*-86-1 (**Figure 1C**
121 **and Table S5**). Their association analysis showed that the 6-bp variation in the SSR
122 region was co-segregated with a fiber strength trait under different field conditions in

123 multiple years at five different locations, including Shihezi, Xinjiang province, the most
124 important cotton production area in China (**Figure S2B**).

125 The haplotype association studies of 268 cotton cultivars (Fang et al., 2017) with
126 marker K222 also showed an extremely significant association with a fiber strength
127 trait at Korla, Xinjiang province (p -value = $3.13e-4$, Student's test), Nanjing, Jiangsu
128 province, and Anyang, Henan province, three cotton-growing regions in China
129 (**Figures 1D and S2C**). The accessions with Prema-type *GhUBX* haplotypes had
130 stronger fiber strength than that of the 86-1 type accessions. Of the examined accessions,
131 87.7% (n=235) had the 86-1 genotype and only 12.3% (n=33) contained the Prema
132 genotype (**Table S6**). This suggests that this elite allelic variation have a huge potential
133 for improving the fiber strength in cotton breeding in the future.

134 The full CDS length of the *GhUBX* was 1416 bp in the 86-1 type and 1410 bp in the
135 Prema type (**Figure S2D**). The protein contained an ubiquitin-associated (UBA)
136 domain in the N-terminal, an ubiquitin-associating (UAS) domain in the middle and an
137 ubiquitin regulatory X (UBX) domain in the C-terminal, consistent to other UBX family
138 proteins (Deruyffelaere et al., 2018). The SSR repeat region containing the 6-bp InDel
139 mentioned above was not located within any known domains (**Figure S2E**). *GhUBX*
140 was consistently expressed in all the tested tissues, including roots, stems, leaves and
141 fibers at different developmental stages, and its expression was significantly lower in
142 the fibers of Prema than those of 86-1 (**Figure S2F**).

143

144 ***GhUBX* is associated with the fiber helix**

145 To gain further insight into the biological role of *GhUBX* in determining the fiber
146 strength, we developed a total of 12 independent *GhUBX*-overexpression (OE)
147 transgenic lines through the *Agrobacterium*-mediated transformation method (Wu et al.,
148 2008). According to PCR detection and expression level of *GhUBX* by qRT-pCR, four
149 *GhUBX*-OE lines (designated as OE-120, OE-141, OE-145 and OE-153) with a
150 noticeably increased expression level of the *GhUBX* were selected, and successively
151 self-pollinated to develop pure lines for further analysis (**Figures 2A and S3A**). The

152 fiber strength parameter (cN/tex: centi-Newton per tex) was increased in the transgenic
153 *GhUBX*-OE lines by 24.0%, compared with the receptor W0 (**Figure 2B and Table S7**).
154 Correspondingly, we examined the thickness of cell wall and the number of helix per
155 unit length in the transformed and untransformed cotton lines. Compared with the
156 transgenic receptor W0, the fiber helix number per unit length was increased by 93.2%,
157 the secondary cell wall (SCW) was reduced by 11.71% to 23.83% in the *GhUBX*-OE
158 lines (**Figures 2C-2F, S3B and S3C**). The variation trend of the helix number per unit
159 length was consistent with fiber strength. Therefore, we theorized that the fiber helices
160 might be closely related to the fiber strength. Moreover, both right- and left-handed
161 helices were presented in mature fibers (**Figure 2C**), indicating that handedness was a
162 randomly selected trait. This phenomena in natural cotton cultivar population were
163 further validated. We selected 60 cultivars with extremely higher and lower fiber
164 strength from the worldwide cotton cultivars to measure the number of fiber helix. A
165 positive relationship between fiber helix and strength was found, and the cultivars with
166 higher fiber strength contained more helices (**Figure S4 and Table S8**). These results
167 suggested incensement of the helix number per unit could enhance fiber elasticity,
168 thereby improving the resistance ability of fibers against external pulling forces and
169 stress (Hsieh, 1999; Haigler, 2010).

170

171 **Short tandem repeat variations influence the interaction of GhUBX with GhSPL1**

172 To address the molecular basis of fiber helix regulation by *GhUBX*, yeast two-hybrid
173 (Y2H) assay was performed to identify its interacting proteins (**Table S9**). We detected
174 a strong interaction of GhUBX with ubiquitin 10 and SPIRAL1-like1. It has been
175 reported that UBX is involved in protein degradation, so its interaction with ubiquitin
176 10 is within our expectation. Therefore, we paid more attention to another interacting
177 protein, SPIRAL1-like1. *SPIRAL1-like1* encodes a six-member family gene, sharing an
178 amino acid homology of 60% with the *SPIRAL1* gene in *Arabidopsis*, so, we named it
179 *GhSPIRAL1-like1* (*GhSPL1*). As previously reported (Sedbrook et al., 2004; Nakajima
180 et al., 2004; Wang et al., 2011; Shoji et al., 2004), the *SPIRAL1* gene functions as a

181 plant-specific microtubule-associated protein (MAP) and regulates the helical growth
182 of roots. The interaction between GhUBX and GhSPL1 was double-checked by co-
183 immunoprecipitation (co-IP) and pull-down assays *in vitro* (**Figures 3A and 3B**), by
184 bimolecular fluorescence complementation (BiFC) assays (Qin et al., 2019), and firefly
185 luciferase complementation imaging (LCI) in *planta* (**Figures 3C and 3D**). Strong
186 fluorescence signals in the cortical microtubules were observed in *Arabidopsis*
187 protoplasts and *Nicotiana benthamiana* leaves, indicating that GhUBX and GhSPL1
188 were co-localized with the microtubule marker MAP-65 (**Figures S5C and S5D**). Their
189 subcellular location inferred that GhUBX interacted with GhSPL1 directly in the
190 cortical microtubules of the developing fibers.

191 To test the exact binding region between GhUBX and GhSPL1, GhUBX was divided
192 into several parts to perform the Y2H assay and LCI assay. The Y2H results showed
193 that GhUBX (71 to 153) was responsible for its interaction with GhSPL1 (**Figures 3E**
194 **and 3F**). Further Y2H and LCI assays revealed that the N-terminal amino acids from
195 the 71st to the 81st regions (in Prema) or to the 83rd region (in 86-1) within the SSRs
196 in GhUBX contributed to its interaction with GhSPL1 (**Figures 3D and 3F**). Overall,
197 these results indicated that this SSR region was the key site bonding with GhSPL1, so
198 we named it as GhSPL1 Interacting Site (**SIS**) domain. In addition to the UBA, UBX,
199 and UAS domains previously reported (Deruyffelaere et al., 2018), a novel SIS domain
200 bound by SPIRAL1-like1 is identified in the *GhUBX* gene.

201 In *Arabidopsis*, *SPIRAL1*(*SPR1*) was reported to be involved in twisted growth, and
202 loss of *SPR1* function conferred right-handed twisting of the plant axis, including the
203 roots, stems, and leaves (Sedbrook et al., 2004; Nakajima et al., 2004). There are two
204 *SPIRAL1-like1* orthologs in Upland cotton, *GH_D03G1112* (*GhSPL1*) and
205 *GH_A03G0847* (*GhSPL1-A*). To explore its orthologous function, virus-induced gene
206 silencing (VIGS) assay was used. Young leaves twisted in the *GhSPL1*-downregulated
207 VIGS cotton plants (**Figures S6A-S6C**). At the boll opening stage, we harvested the
208 mature fiber of *GhSPL1*-silenced group and TRV:00 plants (CK), while, both fiber
209 strength and helix number were significantly enhanced in *GhSPL1*-downregulated

210 plants than CK (**Figures S6D, S6E and Table S10**). Ectopic expression of *GhSPL1* in
211 the *spiral1* mutant lines: CS6546 and SALK_048697, which displayed helical growth
212 phenotypes in the roots of plants (Nakajima et al., 2004), rescued the helical growth of
213 root epidermal cells in *Arabidopsis*, consequently, they became normal, just as the wild
214 type (**Figures S7A-S7C**), that further confirmed the orthologous role of this gene in
215 regulating twisted growth in plants.

216

217 **GhUBX could bridge ubiquitin to GhSPL1 via its UBA domain and SIS domain**

218 Several UBX-containing proteins form a bridge for ubiquitin with substrate proteins in
219 yeast (Lee et al., 2017; Neuber et al., 2005; Schuberth and Buchberger, 2005). We
220 identified the same interaction of GhUBX with the conserved ubiquitin and its UBA
221 domain by Y2H assay (**Figures S8A and S8B**). Through the combination of a prey
222 constructed expressing ubiquitin, a bait constructed expressing GhSPL1 and
223 sectionalized GhUBX bridge proteins (**Figure 4A**), the interactions among GhUBX,
224 GhSPL1, and ubiquitin were detected by Y3H assay. Yeast cells co-transformed with
225 pGADT7-ubiquitin and pBridge-P, pBridge-8, pBridge-NP, pBridge-N8, and pBridge-
226 C grew in the SD/-Leu-Trp medium but not in the SD/-Leu-Trp-His-Ade medium
227 (**Figure 4B**), indicating that ubiquitin did not interact with GhSPL1. When the
228 mentioned yeast cells were spotted on the SD/-Leu-Trp-His-Met medium, they could
229 grow well, except for pGADT7-ubiquitin and pBridge-C (**Figure 4B**), indicating that
230 GhUBX is the bridge between GhSPL1 and ubiquitin and the SIS domain in the N-
231 terminal of GhUBX acts as a key player in the binding of GhUBX to GhSPL1.

232

233 **GhUBX could degrade GhSPL1 via the ubiquitin 26S-proteasome pathway**

234 *In vitro* assays using purified GST-GhUBX and His-GhSPL1 proteins, GhSPL1 could
235 be ubiquitinated in the presence of E1, E2, and GhUBX proteins (**Figure 4C**), the
236 ubiquitination of GhSPL1 by GhUBX promoted its complex degradation. Also,
237 GhSPL1 was degraded by 26S-proteasome pathway. When GhUBX and GhSPL1 both
238 existed with MG132 (a 26S proteasome-specific inhibitor), the signals could be

239 detected with antibody-RFP and antibody-GhSPL1 due to the degradation mediated by
240 GhUBX could be suppressed by MG132 (**Figure 4D**). The more GhUBX led to the
241 decreased of GhSPL1 in developing fiber of Prema, which was opposite in 86-1
242 (**Figures S8C-S8E**). This degradation was further proved by semi-*in vivo* degradation
243 assays that GhUBX-Prema could degrade GhSPL1 faster than GhUBX-86-1 (**Figure**
244 **4E**). Furthermore, the overexpression of *GhUBX* significantly decreased the content of
245 GhSPL1 in developing fibers (**Figure S9**). All these results revealed that GhUBX could
246 regulate the fiber helix in a complex regulatory pathway. In this complex, GhUBX,
247 serving as a functional E₃ ligase, makes effect as the bridge between ubiquitin and
248 GhSPL1 *via* its UBA and SIS domains, and the GhSPL1 is ubiquitinated and further
249 degraded *via* the ubiquitin 26S-proteasome-dependent pathway (**Figure 5**). The
250 reduced GhSPL1 contents lead to the increment of the fiber helices by regulating the
251 stability of the microtubules and, hence, influencing the fiber strength of the transgenic
252 cotton.

253

254 **Discussion**

255 ***GhUBX* regulates cotton fiber strength by helix**

256 Fiber strength is attributed to the rigidity of the cellulosic chains, the frequency and
257 distribution of the reversal, convolitional structure of the microfibril helix and
258 orientation and other characteristics (Hsieh, 1999). Until now, there have been many
259 studies on the influence of SCW on fiber strength. A large number of genes, such as
260 *TALE* superfamily genes, *GhXLIM6*, *GhKNL1*, and *GhCesA4*, and transcription factors,
261 including *MYB*, *NAC*, and *GhTCP4*, have been reported as probably being involved in
262 SCW thickening (Huang et al., 2019; Zhang et al., 2018; Sun et al., 2020; Ma et al.,
263 2019; Li et al., 2018; Gong et al., 2014; Cao et al., 2020). However, the fiber strength
264 is a comprehensive trait not only depending on the thickness of SCW, but also
265 convolitional structure of the microfibril helix and orientation and other characteristics
266 (Hsieh, 1999). The ubiquitin-proteasome system has been explored extensively
267 including autophagy, nuclear transport of specific proteins, repair of DNA, and a

268 multitude of signal transduction pathways (Varshavsky, 2017). Protein with UBX
269 domain, serving as a functional E3 ligase in ubiquitin–proteasome system, participating
270 in lipid droplet formation and degradation of lipid droplet proteins (Wang et al., 2011;
271 Zhang et al., 2017; Feng et al., 2018; Kretzschmar et al., 2018). While, the role of UBX
272 protein in controlling fiber strength has not yet been elucidated. In this study, one new
273 gene containing an UBX domain was identified in two cotton cultivars, Prema and 86-
274 1 (**Figure 1**). The role of GhUBX, a functional E₃ ligase, has been suggested in the
275 ubiquitination and degradation of GhSPL1 *via* the ubiquitin 26S-proteasome dependent
276 pathway (**Figure 4**). The N-terminal of GhUBX has a short tandem repeat variation as
277 the SIS domain, which plays a vital role in binding protein GhSPL1 (**Figure 3**). The
278 GhUBX-OE lines have less GhSPL1 content in the developing fiber cell, leading to the
279 enhancement of the fiber helix compared with the wild type (**Figures S6 and S9**). In
280 general, *GhUBX*-OE lines have thinner SCW than W0, whereas the improvement of
281 fiber strength in the *GhUBX*-OE lines is contributed mostly by more helices. It is
282 amazing, but it makes sense. As for the helices, it is easy to imagine that more helices
283 could enhance the fiber strength (Martinez-Sanz et al., 2017). More helix enhance fiber
284 elasticity due to the improved resistance to external pulling forces, the effects of helix
285 rescue the adverse effect of the reduction of SCW.

286 SPIRAL1 is one of the first proteins identified that determines the twisted growth of
287 plants (Sedbrook et al., 2004; Nakajima et al., 2004). It is a key dynamic microtubule-
288 based modulator for helical growth (Galva et al., 2014). Recent research has confirmed
289 that SPIRAL1 could bind to microtubules and regulate their stability, the spatial
290 distribution of cell wall microfibrils could influence the polarity of cell expansion
291 (Wang et al., 2011). Membrane-assisted cortical microtubules guide the arrangement of
292 cellulose microfibrils (Paredes et al., 2006), thus effects the cell twisting in fiber
293 development (Sambade et al., 2014). The arrangement of fibrils on the surface of fibers
294 at the secondary synthesis stage could influence fiber quality (Han et al., 2013). Recent
295 research has confirmed the actin and microtubule cooperate to pattern the cell wall and
296 growth (Yanagisawa et al., 2015). The arrangement of fibrils affects not only thickness

297 of SCW, but also helix of fiber. We suggest here that the differential expression of
298 *GhUBX* in fibers at the SCW synthesis stage between two parents finally influences the
299 helicity of mature fibers. The increased helices in fiber can make fiber stronger in
300 GhSPL1 silencing group than CK group (**Figure S6 and Table S10**). The variations in
301 cell wall composition result in variable fiber strengths in cotton. And how the
302 microfiber orientation affects the twist of the fibers will be the focus in our future
303 research. We believe that the arrangement of the cortical microtubules can influence the
304 cellulose alignment and the twisting of mature fibers ensures their strength *via* GhSPL1
305 degradation in developing fiber cells by GhUBX. In the cotton cultivars, the number of
306 helix is positively correlated with fiber strength (**Figure S4**). Further investigation is
307 necessary to explore how GhSPL1 regulates the cortical microtubule alignment or
308 microfiber orientation and influences fiber twisting and SCW thickness.

309

310 **Short tandem repeat variation in *GhUBX* correlates with the fiber strength**

311 SSRs are widespread in eukaryotic genomes as short tandem reiterations of sequence
312 motifs. In recent years, there have been an increasing number of reports on their effects
313 on a variety of complex traits (Hammock and Young, 2005). Many debilitating diseases
314 are caused by the repeat expansions in noncoding regions of their resident genes
315 including Huntington disease and hereditary ataxias (Fondon and Garner, 2004; Gatchel
316 and Zoghbi, 2005; Mirkin, 2007; Fotsing et al., 2019). In plants, SSRs are widely used
317 in marker-assisted breeding. The roles of the number of SSR units in regulating
318 phenotypes have been reported in multiple recent studies. For example, microsatellites
319 in starch-synthesizing genes have been found in relation to starch physicochemical
320 properties in waxy rice and 18 types of microsatellites, SNPs, and sequence-tagged sites
321 at three genes are related to starch synthesis (Bao et al., 2002; Bao et al., 2006). A (CT)_n
322 repeat length variation in 5'-UTR of the *CaIMP* gene might regulate phytic acid levels
323 to confer drought tolerance in natural populations of chickpeas (Joshi-Saha and Reddy,
324 2015). Also, a polymorphic (GA/CT)_n varying motif difference of the tryptophan
325 decarboxylase gene at 5'UTR could influence promoter activity (Kumar and Bhatia,

2016). More recently, a variation of specific microsatellite motif size and type enriched in differentially expressed transcripts among latitudinal populations of the common sunflower is reported (Ranathunge et al., 2018).

In *GhUBX*, there are three conservative domains: UBA-like, UAS, and UBX, and a 6-bp deletion located outside of these three domains (**Figure S2E**). Our present study has identified two classes of multiple microsatellite alleles, based on the nucleotide difference between cotton accessions CTCGGCCTCT(GCCTCC)_{5/6}, which were widely existed throughout the genome of *G. hirsutum*, this was authenticated by the microsatellite genotyping of 268 cultivars and RILs with K222 (**Figures S2B and S2C**). The correlation analysis suggested that *GhUBX* was significantly associated with fiber strength (**Figure S2**). The overexpressed transgenic lines demonstrated a high-fiber strength phenotype (**Figure 2B and Table S7**) and confirmed that *GhUBX* associated with fiber strength significantly.

In summary, we cloned the causal gene, *GhUBX*, underlying the major fiber strength QTL *qFS-D3-1*, through map-based cloning in cotton. GhUBX interacts with GhSPL1, and results in significantly increased fiber strength in transgenic cotton. Moreover, GhUBX can act as a bridge between ubiquitin and GhSPL1 and reduce the GhSPL1 content in fiber cells *via* the ubiquitin 26S-proteasome pathway (**Figure 5**). The illustration of fiber strength formation mechanism could enhance our understanding of fiber development theoretically and lay a foundation for improving fiber quality. How GhUBX regulates SCW directly and/or GhUBX interacts with other unknown proteins to influence SCW remains to be elucidated.

348

349 **Limitations of the study**

To further verify our results in natural populations, we tried to use InDel GWAS analysis to obtain the correlation between 6-bp InDel within GhUBX and fiber strength, no significant peaks for fiber strength were identified on Chr. D03. We also found that genotyping results based on PCR and Sanger sequencing were not consistent with the haplotypes as determined by resequencing data, indicating a high error of genotyping

355 based on low-coverage resequencing data. For now, InDel markers are nearly used in
356 GWAS analysis because of the high prevalence of false positives in InDels identified
357 from low-coverage short-read sequencing.

358

359

360

361

Journal Pre-proof

Main figure titles and legends

362

363 **Figure 1. Cloning of *qFS-D3-1*.** (A) *qFS-D3-1* was mapped on the D3 chromosome between the
 364 k5209 and k5222 markers using an F₂ generation. *qFS-D3-1* was further fine-mapped to a region
 365 between the K5219 and K5221 markers using 1864 individuals. The mapping area was narrowed
 366 down to a 0.93-Mb genomic interval, and *GH_D03G0985* (*GhUBX*) was selected as a major gene.
 367 (B) GhUBX contains 4 exons and 3 introns. The DNA sequence alignment of the *GhUBX* gene
 368 exhibits a 6-bp difference between Prema and 86-1. (C), (D) Genotype association analysis of the
 369 k222 marker for fiber strength in RILs and natural populations. 86-1: 86-1 genotype
 370 (GCCTCT(GCCTCC)₆GTCC), Prema: Prema genotype (GCCTCT(GCCTCC)₅GTCC); p-values
 371 were determined by the Student's t-test (**, $p < 0.01$).

372

373 **Figure 2. Transgenic validation of GhUBX conferring increase of fiber strength.** (A) qRT-
 374 PCR analysis of *GhUBX* expression in fibers of the receptor W0 and the *GhUBX*-OE at 15, 20, and
 375 25 DPA. *Histone3* was used as the internal control. (B) Comparison of fiber strength (cN/tex: centi-
 376 Newton per Tex) parameters between the W0 and transgenic cotton plants for three seasons, with
 377 three replicates for each sample. (C) Scanning electron microscopy (SEM) of W0 and transgenic
 378 mature fibers (OE-120, OE-141, OE-145 and OE-153). Scale bars on the SEM images are 50 μ m.
 379 (D) Measurement of the helix numbers by SEM in the W0 and transgenic cotton lines; total length
 380 of each mature fiber was over 15 cm. $n > 30$. (E) Transmission electron microscopy (TEM) of the
 381 W0 and transgenic mature fibers (OE-120, OE-141, OE-145 and OE-153); scale bars on the TEM
 382 images are 0.5 μ m. (F) Cell wall thickness of mature fiber in the W0 and transgenic lines. The data
 383 in A, B, D, and F are shown as the mean (\pm SD) of three experimental replicates; p-values were
 384 determined by the Student's t-test (*, $p < 0.05$; **, $p < 0.01$).

385

386 **Figure 3. GhUBX interaction with GhSPL1.** (A) GhUBX from Prema and 86-1 can interact with
 387 GhSPL1 in a co-IP assay. Co-IP was carried out with anti-GFP agarose from total isolated proteins,
 388 and immunoblotting analysis was done with anti-GFP and anti-RFP antibodies. (B) The direct
 389 interaction between GhUBX and GhSPL1 tested with the *in vitro* pull-down assay. (C) The
 390 interaction of GhUBX and GhSPL1 confirmed by the BiFC assay in *N. benthamiana*. *MAP-65-RFP*

391 as a marker for microtubule localization was cotransformed. Scale bars = 10 μ m. **(D)** GhUBX (1 to
392 153), but not GhUBX (153 to 472), can interact with GhSPL1 as indicated by the split firefly
393 luciferase complementation imaging assay. **(E)** The domain composition of GhUBX amino acid
394 sequence: UBA (ubiquitin-associated domain); SIS (SPL1 interacting site); UAS (ubiquitin-
395 associating domain); UBX (ubiquitin regulatory X domain). **(F)** Y2H assays to investigate the
396 interaction between different domains of GhUBX and GhSPL1. N-terminal amino acids 71 to 83 of
397 86-1 and 71 to 81 of Prema contribute to the interaction with GhSPL1. Deletion of two amino acids
398 in aligning amino acid sequences caused the differences in interactions between GhUBX and
399 GhSPL1. pGBKT7-*p53* and pGADT7-*T-antigen* were used as positive controls, and pGBKT7-
400 *Lamin c* and pGADT7-*T-antigen* as negative controls.

401

402

403 **Figure 4. Degradation of a complex ubiquitin/UBX/SPL1 molecule via ubiquitin 26S-**
404 **proteasome pathway. (A)** The constructs expressing both the bait and bridge proteins for Y3H
405 assays. **(B)** GhUBX acts as a bridge between ubiquitin and GhSPL1 in Y3H interactions. Ubiquitin
406 did not interact with GhSPL1 directly, but GhUBX (UBA and SIS domain), ubiquitin and GhSPL1
407 showed interactions. **(C)** GhUBX ubiquitinates GhSPL1 *in vitro*: ubiquitinated GhSPL1-His was
408 detected by anti-ubiquitin and anti-His. * indicates ubiquitinated GhSPL1. **(D)** GhUBX promotes
409 the degradation of GhSPL1 via the 26S-proteasome *in vivo*. Immunoblotting analysis of protein
410 extracts corresponding to agroinfiltrated *N. benthamiana* leaves with the indicated plasmids in the
411 presence or absence of MG132. Immunoblotting analysis was done with anti-GFP and anti-RFP
412 antibodies. The anti-RFP antibody served as a loading control. The mRNA expression levels of
413 GhUBX and GhSPL1 were analyzed by RT-PCR, and the mRNA expression level of *EF1 α* was used
414 as the internal control. **(E)** GhUBX promotes the degradation of GhSPL1 in a semi-*in vivo* protein
415 degradation assay. Extracts from GhSPL1-RFP were mixed together with GhUBX-Prema-GFP,
416 GhUBX-86-1-GFP or GFP-vector, after which the mixtures were treated with CHX and ATP in the
417 presence of MG132 or DMSO for different times. GFP antibody and RFP antibody were used for
418 detecting the abundance GhUBX and GhSPL1. Rubisco was used as a loading control.

419

420

421 **Figure 5. A working model of GhUBX ubiquitinates GhSPL1 via ubiquitin 26S-proteasome**
422 **pathway.** In the working model, GhUBX is a bridge protein that functions by bonding to ubiquitin
423 and GhSPL1 *via* its UBA domain and SIS domain, respectively. GhUBX recruits GhSPL1 for
424 degradation via the ubiquitin 26S-proteasome pathway. In transgenic cotton, overexpression of
425 GhUBX reduces the GhSPL1 protein in microtubule, leading to more helices in mature fibers.

426

427 **Supplemental figure and table titles**

428 **Table S1.** All primers developed and used in the present research. Related to STAR Methods.

429 **Table S5.** Fiber strength and genotype (Prema and 86-1) of 183 recombinant inbred lines. Related
430 to Figures 1 and S2.

431 **Table S6.** Fiber strength (cN/tex) and genotype (Prema and 86-1) of 268 cotton varieties. Related
432 to Figures 1 and S2.

433

434

435 **Acknowledgements**

436 This study was financially supported by grants from the National Natural Science
437 Foundation of China (31822036; 31701469); the National Research and Development
438 Project of Transgenic Crops of China (2016ZX08009-003); the Fundamental Research
439 Funds for the Central Universities (2019XZZX004-03, 2019XZZX004-13); and
440 Leading Innovative and Entrepreneur Team Introduction Program of Zhejiang
441 (2019R01002). We would like to thank the Bio-ultrastructure Analysis Lab. of the
442 Analysis center of Agrobiological and Environmental Sciences, Zhejiang Univ. for the
443 TEM and SEM assays.

444

445

446 **Author contributions**

447 T.Z. and Y.Z. designed the research; Y.Z., C.X., W.S., Y.W., Z.N., S.W., Z.H., Z.S., Y.Z.,
448 and L.F. performed experiments; T.Z., Y.Z., Y.H., and Y.W. analyzed the data; T.Z., and
449 Y.Z. wrote the paper. All authors discussed results and commented on the manuscript.

450

451

452 **Declaration of interests**

453 The authors declare no competing financial interests.

454

455 **STAR Methods**456 **Key resources table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti-His	Sigma-Aldrich	Cat# SAB1305538; RRID: AB_2687993
Anti-GST	Sigma-Aldrich	Cat# G7781, RRID:AB_259965
Anti-GFP	Sigma-Aldrich	Cat # G1546; RRID: AB_1079024
Anti-RFP	Sigma-Aldrich	Cat# AB3528, RRID:AB_91496
Bacterial and Virus Strains		
Trelief™ 5α Chemically Competent Cell (DH5a)	Tsingke	Cat# TSC01
GV3101(pSoup) Chemically Competent Cell	WeidiBio	CAT#: AC1002
EHA105 Electroporation-Competent Cell	WeidiBio	CAT#: AE1010
Chemicals, Peptides, and Recombinant Proteins		
Murashige and Skoog Basal Medium	Sigma-Aldrich	M5519
MG132	Sigma-Aldrich	Cat#M8699-1MG
MgCl ₂ ·6H ₂ O	SIGMA	M2670
MES	SIGMA	M8250
CaCl ₂	SIGMA	C5670

Kanamycin sulfate	Amresco	0408
Rifampicin	SIGMA	R3501
Ampicillin	Amresco	0399
Silwet L-77	GE Healthcare	SL77080596
Dimethyl sulfoxide	SIGMA	D8418
PMSF	American Bioanalytical	AB01620
cOmplete, Protease Inhibitor Cocktail	Sigma-Aldrich	Cat#4693159001
Critical Commercial Assays		
Scanning and transmission electron microscope analysis	Bio-ultrastructure Analysis Laboratory of the Analysis Center, Agrobiology and Environmental Sciences, Zhejiang University	NA
Cotton transformation	Biotechnology Research Center, Shanxi Academy of Agricultural Sciences	NA
Fiber quality measurement	Cotton Quality Supervision, Inspection and Testing Center, Ministry of Agriculture, Xinjiang, China.	NA
Experimental Models: Organisms/Strains		
TM-1	This paper	NA
Prema	This paper	NA
86-1	This paper	NA
W0	This paper	NA
<i>N. benthamiana</i>	This paper	NA
<i>Arabidopsis thaliana</i> ecotypes Col-0 and Ler	This paper	NA

<i>Arabidopsis thaliana: spirall</i>	ABRC	SALK_048697
<i>Arabidopsis thaliana: spirall-1</i>	ABRC	CS6546
Oligonucleotides		
Primers are listed in Table S1	This paper	NA
Recombinant DNA		
35S::GhUBX-86-1-GST	This paper	NA
35S::GhUBX-Prema-GST	This paper	NA
35S::GhSPL1-HIS	This paper	NA
35S::GhSPL1-RFP	This paper	NA
35S::GhUBX-Prema-GFP	This paper	NA
35S::GhUBX-86-1-GFP	This paper	NA
35S::MAP65-RFP	Li et al.,2017	NA
Software and Algorithms		
Prism 6	https://www.graphpad.com/	NA
ImageJ	https://imagej.net/Welcome	NA

457

458 **Resource availability**459 **Lead contact**

460 Further information and requests for resources and reagents should be directed to and
 461 will be fulfilled by the Lead Contact, Tianzhen Zhang (cotton@njau.edu.cn).

462 **Materials availability**

463 All materials are available after completion of the respective material transfer
 464 agreements.

465 **Data and code availability**

466 Any data information reported in this paper is available from the lead contact upon
 467 request.

468

469 This study did not generate new reagents or software code.

470

471 **Experimental model and subject details**

472 Plant materials

473 Upland cotton (*Gossypium hirsutum*) plants, W0, TM-1, Prema, and 86-1, were
474 cultivated in the field at the experimental station of Nanjing Agricultural University
475 (NJAU) in China. The high-fiber strength parental line Prema, developed in California
476 by a cross (AXTE-1 × NM49-2) × (C6TE × NMB3080), is a high-fiber quality Acala
477 cultivar (Ulloa and Jr, 2000). Prema was an introgressed line from *Gossypium thurberi*
478 and *Gossypium barbadense*. The low-fiber strength parental line 86-1 had been widely
479 cultivated in the Yangtze River and Yellow River cotton growing regions since the
480 1980s (Jian et al., 2003). In 2005, Prema and 86-1 were crossed at the Jiangpu Breeding
481 Station, Nanjing Agricultural University (JBS/NAU), and RIL families were
482 constructed and used for the present research (Ning et al., 2014). RILs were planted in
483 China at Jiangpu, Nanjing/Jiangsu, Sanya/Hainan, and Shihezi/Xinjiang in 2009 and in
484 Shihezi/Xinjiang and Dafeng/Jiangsu in 2010. Fiber cells at different developmental
485 stages were carefully removed from seeds and immediately snap-frozen in liquid
486 nitrogen for DNA and RNA extraction.

487

488 Methods details**489 Mapping population development and fiber quality measurement**

490 According to the genotype and location of the QTL mapped on the RILs population
491 (Acala Prema × 86-1), we selected four RILs: RIL168, RIL98, RIL120, RIL43, which
492 with fine fiber strength in diverse environments for years to cross with 86-1 in the
493 summer of 2014 at the JBS/NAU. The F₁ plants were self-pollinated in the winter of
494 2014 in Hainan Island to produce F₂ progeny. A total of 1864 F₂ individuals, including
495 four F₂ populations were planted in the summer of 2015 in JBS/NAU, the sample size
496 of each F₂ populations crossed between RIL168, RIL98, RIL120 and RIL43, and 86-1
497 was 747, 355, 417 and 345 individuals, respectively. All naturally opened bolls were
498 hand-harvested to gain fiber that could be measured for fiber quality, including fiber
499 length (FL, mm), strength (FS, cN/tex), micronaire (FMIC), elongation (FE), and
500 uniformity ratio (FU). Leaf samples were collected from 6-week-old plants of each F₂

501 generation grown in the field.

502 For comparative field trials, the transgenic cotton and WT were planted in a
503 randomized block design with three replications. Each experimental replication
504 included approximately one 4.5-m long row, with 75 cm between rows and 35 cm
505 between plants in each row. Five plants in the middle of each row were tagged for
506 harvesting. The middle cotton bolls were harvested and the fiber quality was
507 investigated. Fiber quality traits included the FL, FS, FMIC, FE, and FU. The testing
508 standard was the GB/T 20392-2006 HVI Cotton Fiber Physical Properties Test Method.
509 Fiber samples were measured by the Cotton Quality Supervision, Inspection and
510 Testing Center, Ministry of Agriculture, Xinjiang, China.

511

512 **Marker analysis and genetic map construction**

513 Cotton genomic DNA was extracted from young leaves using a modified cetyl
514 trimethylammonium bromide method (Paterson et al. 1993). To enrich markers within
515 the QTL region, SNP markers identified by restriction site-associated DNA sequencing
516 and InDel markers identified by transcriptome sequencing were used for fine mapping
517 (Wang et al., 2015). SSR and InDel markers were developed based on the resequencing
518 data between 86-1 and Prema. The markers used in the present research are listed in
519 **Table S1**. Primers were synthesized by TSINGKE (Beijing, China) and mapped on Chr.
520 D03, which displayed a clear polymorphism between two parents. Joinmap3.0 (Ooijen,
521 2001) was used to construct the genetic map of the primary QTL region on Chr. D03,
522 with a LOD score of 6.0 and a recombination frequency of 0.40. The complex interval
523 mapping method of Windows QTL Cartographer 2.5 (Basten, 2001) was used to
524 identify the QTL for the fiber quality traits. QTLs were determined to be significant if
525 the corresponding likelihood ratio score was greater than 11.5. MapChart2.2 (Voorrips,
526 2002) was used to develop the linkage group and anchor the QTL. The QTL was named
527 starting with q, followed by an abbreviation of the trait name, the name of the
528 chromosome, and the QTL number affecting the trait on the chromosome (QTL + trait
529 + number) (McCouch, 1997).

530

531 **The quantitative RT-PCR analysis**

532 The qRT-PCR specific primers were designed using Integrated DNA Technologies
533 online (<https://sg.idtdna.com>). *Histone3* (AF024716) was used as an internal control.
534 qRT-PCR was carried out on an Applied Biosystems 7500 Fast Real-Time PCR System
535 (Life Technologies, Foster City, CA, USA) in a 20- μ l volume containing 100 ng of
536 cDNA, 4 pM of each primer, and 10 μ l of AceQ qPCR SYBR Green Master Mix
537 (Vazyme, Nanjing, China) according to the manufacturer's protocol. The data were
538 evaluated using the comparative cycle threshold method described by Livak and
539 Schmittgen (Livak and Schmittgen, 2001). Three biological replicates (three samples
540 harvested from three plants, one from each) were performed per reaction, each with two
541 technical replicates (using the same sample). Mean values and standard errors were
542 calculated according to the data from three replicates.

543

544 **Scanning and transmission electron microscope analysis**

545 Samples of mature fibers from Prema, 86-1, transgenic cotton lines, and W0 were
546 dehydrated, fixed on the observation table, sputter-coated with silver using an E-
547 1010/E-1020 ion sputter (Hitachi, Japan), and imaged using an SU8010 scanning
548 electron microscope at 3.0 kV (Hitachi, Japan).

549 The mature fibers of the transgenic cotton were observed using a transmission
550 electron microscope. Transverse sections of the fiber samples were fixed in 2.5%
551 glutaraldehyde in a phosphate buffer overnight at 4°C and then in 1% OsO₄ for 2 hours.
552 The samples were further dehydrated through the application of step-graded ethanol
553 and embedded in Spurr's medium prior to ultrathin sectioning. Sections (90 nm thick)
554 were cut with an ultramicrotome (EMUC7, 645 Leica, Germany) and collected on
555 nickel mesh. The sections were air dried, stained, and viewed with a Hitachi H-7650
556 TEM at 80 kV (Bio-ultrastructure Analysis Laboratory of the Analysis Center of
557 Agrobiological and Environmental Sciences, Zhejiang University). Five to 10 nonserial
558 sections per genotype from fibers were examined per line. At least 10 cell wall

559 interfaces between the fiber and fiber base from fibers per genotype were examined.
560 The same mature fiber samples used for fiber quality investigation were also used for
561 scanning and transmission electron microscopic analysis.

562

563 **Cotton transformation**

564 For the overexpression constructs, primers with added *Bam*H I and *Sma* I, were used to
565 amplify the open reading frame of GhUBX from 86-1, which was then cloned into the
566 pBI121 vector under the control of the constitutive cauliflower mosaic virus 35S
567 promoter. GhUBX-overexpression (GhUBX-OE) construct was introduced into *G.*
568 *hirsutum* acc. W0 via *Agrobacterium tumefaciens*-mediated transformation.
569 *Agrobacterium* strain LBA4404 holding the pBI121 plasmid vector was grown in
570 Luria–Bertani liquid medium supplemented with 50 mg/L kanamycin and 10 mg/L
571 rifampicin at 28°C for 24 h. The bacteria were resuspended in liquid MSB₁ medium
572 and the standard OD600 was adjusted to 0.3~0.5. Embryogenic calli were inoculated
573 with the *Agrobacterium* suspension for 20 min and subsequently blotted dry with sterile
574 filter papers. Then, the calli were dispersed and cocultured on MSB₁ medium placed on
575 filter paper in the dark (Wu et al., 2008). The homozygosity of the transgenic plants
576 was determined using the kanamycin selection marker coupled with PCR-based
577 genotyping. DNA polymerase (I5™ 2 × High-Fidelity Master Mix) was purchased from
578 TSINGKE (Beijing, China). The primers used for vector construction and PCR-based
579 screening were listed in **Table S1**.

580

581 **Subcellular localization**

582 For subcellular localization of *GhUBX* in *N. benthamiana* leaves, the coding sequence
583 of GhUBX was cloned into the vector pBinGFP4 to form GhUBX-GFP. *A. tumefaciens*
584 strain GV3101 carrying the construct was used to infiltrate 6-week-old *N. benthamiana*
585 leaves. For the subcellular location of *GhUBX* and *MAP-65* in *Arabidopsis* protoplast,
586 the *GhUBX*-GFP4 and *MAP-65* plasmid were transiently transformed *via* PEG/Ca²⁺ and
587 conducted as previously reported (Confraria and Baena-González, 2016). *MAP-65* used

588 as microtubule marker (Lucas and Shaw, 2012; Li et al., 2017). Analysis was performed
589 on a Zeiss LSM780 confocal microscope using a 488-nm excitation laser for GFP and
590 a 561-nm laser for RFP. Images were processed using Zen 2009 software.

591

592 **Yeast two-hybrid assays and yeast three-hybrid assays**

593 The yeast two-hybrid (Y2H) assay was performed using the Gal4 vector system
594 (Clontech, USA). Sectionalized GhUBX and GhSPL1 were cloned into both the
595 pGBKT7 vector and pGADT7 vector. The coding regions of GhUBX (86-1-1-472),
596 GhUBX (Prema-1-470), GhUBX (153-472) (non-distinctive region), GhUBX (86-1-1-
597 83), GhUBX (Prema-1-81), GhUBX (86-1-71-153), GhUBX (Prema-71-153), GhUBX
598 (1-50) (non-distinctive region), and GhSPL1 were cloned into both the pGBKT7 vector
599 and pGADT7 vector, respectively. The constructs were cotransformed into yeast strain
600 Y2H. The transformed cells were adjusted to $OD_{600} = 0.4 \sim 0.6$ and grown in SD/-Trp-
601 Leu or SD/-Trp-Leu-His-Ade plates for 3 to 7 days at 30°C.

602 Y3H assay was performed based on the pBridge vector system (Takara Bio, Japan).
603 The conserved ubiquitin sequence was fused to GAL4 AD in pGADT7. To construct
604 the pBridge-GhSPL1-GhUBX, the full-length coding sequence of the GhSPL1 coding
605 region was cloned into multiple cloning site (MCS) I of the pBridge vector fused to the
606 GAL4 BD domain, and the sectionalized GhUBX coding region was cloned into MCS
607 II of the pBridge vector. The transformed cells were grown in SD/-Trp-Leu-Met plates
608 and SD/-Trp-Leu-His-Met plates for 5 to 8 days at 30°C.

609

610 **Pull-down assays**

611 The coding sequences of GhUBX-Prema/86-1 and GhSPL1 were cloned into the
612 pGEX-4T-1 vector and pET-32a vector to generate the constructs to express GST-
613 GhUBX-Prema/86-1 and His-GhSPL1, respectively. The pull-down assays were
614 performed in accordance with previous reported methods (Xia et al., 2013).

615

616 **GhUBX polyclonal antibody preparation**

617 For the preparation of the GhUBX polyclonal antibody, a 1416-bp coding-region
618 fragment encoding a 472–amino acid peptide of GhUBX-86-1 was cloned into a
619 pET32a vector (AOGENE, Nanjing, China). The recombinant protein was expressed in
620 *Escherichia coli* DE3 (Transgen, Beijing, China) and purified to produce rabbit
621 polyclonal antibodies (prepared by AOGENE of China).

622

623 **Co-immunoprecipitation assays**

624 Co-immunoprecipitation (co-IP) was conducted as previously described (Liu et al.,
625 2010). In brief, *GhSPL1*-RFP and *GhUBX*-GFP or vector-GFP and vector-RFP were
626 transiently co-expressed in *N. benthamiana* leaves for 3 days. At 12 hours before
627 sample collection, 100 μ M of the 26S-proteasome inhibitor MG132 was infiltrated.
628 Lysates were incubated with anti-GFP or anti-RFP affinity M2 beads (Sigma-Aldrich,
629 St. Louis, MO, USA) at 4°C for 2 hours. The beads were washed three times with PBS,
630 and the immunoprecipitated proteins were examined by immunoblotting (Ruiyuan
631 Biotech, Nanjing, China).

632

633 **Luciferase imaging assays**

634 For the firefly LUC complementation imaging assays, the sectionalized sequences of
635 GhUBX-P, GhUBX-8, GhUBX-P-1-153, GhUBX-8-1-153, and GhUBX-153-472 were
636 respectively ligated with the N-terminal fragment of luciferase (nLUC) to form
637 sectionalized-GhUBX-nLUC. The full-length coding sequences of GhSPL1 were fused
638 with the C-terminal fragment of luciferase (cLUC). FT1 and FD worked as positive
639 controls (Qin et al., 2019). The images were taken by a low-light, cooled, charge-
640 coupled device imaging apparatus (Tanon, Fremont, CA, USA) (Kong et al., 2015).

641

642 **BiFC assays**

643 For the BiFC assays, sectionalized sequences of *GhUBX* were cloned into YNE and
644 fused with the N-terminus of YFP; and *GhSP11* was cloned into YCE and fused with
645 the C- terminus of YFP. The recombinant constructs were co-transformed in pairs into

646 young *N. benthamiana* leaves, and *MAP65-RFP*, which served as a marker for the
647 microtubule localization, was also co-transformed (Lucas and Shaw, 2012; Li et al.,
648 2017; Burkart and Dixit, 2019). The fluorescence signal was observed using a Zeiss
649 LSM780 confocal microscope.

650

651 **Virus-induced gene silencing (VIGS) assay**

652 For Virus-induced gene silencing (VIGS) assay, a 296-bp fragment of GhSPL1 cDNA,
653 corresponding to bases 22 to 317 of the GhSPIRAL1-like1 gene, was amplified by PCR.
654 The resulting PCR product was cloned into pTRV2 to produce a vector referred to as
655 pTRV2-GhSPL1. *Agrobacterium* cells respectively carrying pTRV1 and pTRV2-
656 GhSPL1 were re-suspended in an infiltration medium (10 mM MgCl₂, 10 mM MES,
657 200 μM acetosyringone) and adjusted to an OD₆₀₀ of 1.0. The *Agrobacterium* strains
658 containing the pTRV1 and pTRV2-GhSPL1 vectors were mixed at a ratio of 1:1. The
659 *Agrobacterium* suspension was injected into the cotyledons of 10-day-old seedlings,
660 which were placed in the dark for 24 hours and then incubated at 23°C with a 16-hour
661 light/8-hour dark cycle. The TM-1 was used as a receptor to carry on the VIGS assay
662 to silence the *GhSPL1* gene. Empty-vector (TRV: 00) transformed plants were used as
663 experimental controls. The chloroplast alterados 1 (*CLAI*) gene was used as an
664 indicating gene to control the silencing effect (Gao et al., 2011). Thirty seedlings were
665 used as experimental groups to silence the *GhSPL1* gene. The photos were taken three
666 weeks after injection, and leaves were collected for expression detection.

667

668 ***In vitro* ubiquitination**

669 GST-GhUBX-Prema/86-1 and His-GhSPL1 proteins were purified from the *E. coli*
670 strain Arctic-Express. *In vitro* ubiquitination assay was conducted as previously
671 reported (Yang et al., 2015).

672

673 ***In vitro*, *in vivo*, and semi-*in vivo* protein degradation assays**

674 *In vitro* degradation assays were performed in accordance with previously reported

675 methods (Yang et al., 2015). For semi-*in vivo* protein degradation analysis, a GhUBX-
676 Prema-GFP sample, GhUBX-86-1-GFP sample, GhSPL1-RFP sample, and GFP
677 sample were harvested at 3 days after inoculation. The four samples were separately
678 extracted with native extraction buffer (50 mM Tris-MES [pH 8.0], 0.5 M sucrose, 1
679 mM MgCl₂, 10 mM EDTA, 5 mM DTT, and a protease inhibitor cocktail
680 (CompleteMini tablets; Roche, Indianapolis, IN, USA); the indicated concentrations of
681 ATP, CHX, and MG132 or DMSO (control) were added to the protein extract. The
682 mixtures were incubated at 25°C. Samples were removed at different time points for
683 immunoblot analyses.

684 For *in vivo* degradation assays, we coinfiltrated the *Agrobacterium* strains carrying
685 the *GhUBX* (Prema/86-1)-GFP and *GhSPL1*-RFP plasmids into *N. benthamiana* leaves
686 for 3 days. The corresponding empty vectors were used as the controls, and the RFP
687 plasmid was added as an internal control. For proteasome inhibition, leaves were
688 infiltrated with a 10 mM MgCl₂ and 100 mM MG132 solution for 14 hours before
689 sample collection. Samples were collected for protein and RNA extraction. For protein-
690 level analysis, the extracts were analyzed using anti-GFP antibody (1:1000 dilution),
691 anti-RFP antibody (1:1000 dilution), anti-GhUBX antibody (1:1000 dilution), and anti-
692 GhSPL1 antibody (1:1000 dilution). For RNA level expression analysis, RT-PCR was
693 performed.

694

695 **Association mapping**

696 A haplotype association panel composed of 268 diverse cotton cultivars was used in the
697 association analysis of the tandem repeats for GhUBX (**Table S6**). This association
698 population was previously phenotyped for all fiber quality traits (Fang et al., 2017). The
699 significance of difference was analyzed with two-tailed test. Another population
700 composed of 486 cotton cultivars were used to analyze the association between fiber
701 strength and helices number. 20 individuals with highest FS (cN/tex > 35) and 20
702 individuals with lowest FS (cN/tex < 26) were selected from this population and
703 calculated their helices. The helices number and other fiber quality traits are shown in

704 **Table S8.**

705

706 **Quantification and statistical analysis**

707 In this study, significant differences between two samples were determined with Two-
708 tailed paired Student's t test. Error bars represent standard deviation of mean, 'n'
709 represents the sample size, as mentioned in the figure legends. And asterisks indicate
710 the statistical significance: *, $p < 0.05$; **, $p < 0.01$. At least three biological replicates
711 were included. Statistical analysis was performed by GraphPad Prism 6.0. Thickness of
712 cell wall was analysis by ImageJ.

713

714

715

716

717 **REFERENCES**

718

- 719 Bao, S., Corke, H., and Sun, M. (2006). Microsatellites, single nucleotide polymorphisms and a sequence
720 tagged site in starch-synthesizing genes in relation to starch physicochemical properties in nonwaxy rice
721 (*Oryza sativa* L.). *Theor. Appl. Genet.* *113*, 1185-1196.
- 722 Bao, S., Corke, H., and Sun, M. (2002). Microsatellites in starch-synthesizing genes in relation to
723 starch physicochemical properties in waxy rice (*Oryza sativa* L.). *Theor. Appl. Genet.* *105*, 898-905.
- 724 Baskin, T.I. (2005). Anisotropic expansion of the plant cell wall. *Annu. Rev. Cell Dev. Biol.* *21*, 203-222.
- 725 Basten, C.J., Weir, B. S. and Zeng, Z. B. (2001). QTL cartographer, version 1.15. Department of Statistics,
726 North Carolina State University. Raleigh.
- 727 Burkart, G.M., and Dixit, R. (2019). Microtubule bundling by MAP65-1 protects against severing by
728 inhibiting the binding of katanin. *Mol. Biol. Cell* *30*, 1587-1597.
- 729 Cao, J.F., Zhao, B., Huang, C.C., Chen, Z.W., Zhao, T., Liu, H.R., Hu, G.J., Shangguan, X.X., Shan,
730 C.M., Wang, L.J., et al. (2020). The miR319-targeted GhTCP4 promotes the transition from cell
731 elongation to wall thickening in cotton fiber. *Mol. Plant* *13*, 1063-1077.
- 732 Confraria, A., and Baena-González, E. (2016). Using *Arabidopsis* protoplasts to study cellular responses
733 to environmental stress. *Methods Mol. Biol.* *1398*, 247-69.
- 734 Deruyffelaere, C., Purkrtova, Z., Bouchez, I., Collet, B., Cacas, J.L., Chardot, T., Gallois, J.L., and
735 D'Andrea, S. (2018). PUX10 is a CDC48A adaptor protein that regulates the extraction of ubiquitinated
736 oleosins from seed lipid droplets in *Arabidopsis*. *Plant Cell* *30*, 2116-2136.
- 737 Dumais, J., and Forterre, Y. (2012). "Vegetable Dynamicks": The role of water in plant movements. *Annu.*
738 *Rev. Fluid Mech.* *44*, 453-478.
- 739 Fang, L., Wang, Q., Hu, Y., Jia, Y., Chen, J., Liu, B., Zhang, Z., Guan, X., Chen, S., Zhou, B., et al. (2017).
740 Genomic analyses in cotton identify signatures of selection and loci associated with fiber quality and
741 yield traits. *Nat. Genet.* *49*, 1089-1098.
- 742 Feng, H., Li, X., Chen, H., Deng, J., Zhang, C., Liu, J., Wang, T., Zhang, X., and Dong, J. (2018).
743 GhHUB2, a ubiquitin ligase, is involved in cotton fiber development *via* the ubiquitin-26S proteasome
744 pathway. *J. Exp. Bot.* *69*, 5059-5075.

- 745 Fondon, J.W., 3rd, and Garner, H.R. (2004). Molecular origins of rapid and continuous morphological
746 evolution. *Proc. Natl. Acad. Sci. USA* *101*, 18058-18063.
- 747 Fotsing, S.F., Margoliash, J., Wang, C., Saini, S., Yanicky, R., Shleizer-Burko, S., Goren, A., and Gymrek,
748 M. (2019). The impact of short tandem repeat variation on gene expression. *Nat. Genet.* *51*, 1652-1659.
- 749 Galva, C., Kirik, V., Lindeboom, J.J., Kaloriti, D., Rancour, D.M., Hussey, P.J., Bednarek, S.Y., Ehrhardt,
750 D.W., and Sedbrook, J.C. (2014). The microtubule plus-end tracking proteins SPR1 and EB1b interact
751 to maintain polar cell elongation and directional organ growth in *Arabidopsis*. *Plant Cell* *26*, 4409-4425.
- 752 Gao, X., Britt, R.C., Jr., Shan, L., and He, P. (2011). *Agrobacterium*-mediated virus-induced gene
753 silencing assay in cotton. *J. Vis. Exp.* *54*, e2938-e2938.
- 754 Gatchel, J.R., and Zoghbi, H.Y. (2005). Diseases of unstable repeat expansion: mechanisms and common
755 principles. *Nat. Rev. Genet.* *6*, 743-755.
- 756 Gong, S.Y., Huang, G.Q., Sun, X., Qin, L.X., Li, Y., Zhou, L., and Li, X.B. (2014). Cotton KNL1,
757 encoding a class II KNOX transcription factor, is involved in regulation of fibre development. *J. Exp.*
758 *Bot.* *65*, 4133-4147.
- 759 Haigler, C.H. (2010). Physiological and anatomical factors determining fiber structure and utility. In
760 *Physiology of Cotton*, J.McD. Stewart et al., eds. (Springer Science+Business Media), Chapter 4, 33-
761 47.
- 762 Hammock, E.A., and Young, L.J. (2005). Microsatellite instability generates diversity in brain and
763 sociobehavioral traits. *Science* *308*, 1630-1634.
- 764 Han, L.B., Li, Y.B., Wang, H.Y., Wu, X.M., Li, C.L., Luo, M., Wu, S.J., Kong, Z.S., Pei, Y., Jiao, G.L.,
765 et al. (2013). The dual functions of WLIM1a in cell elongation and secondary wall formation in
766 developing cotton fibers. *Plant Cell* *25*, 4421-4438.
- 767 Hsieh, Y.L. (1999). Structural development of cotton fibers and linkages to fiber quality. In *Cotton Fibers*,
768 A.S. Basra, ed. (Binghampton: Food Products Press), 137-166.
- 769 Hu, Y., Chen, J., Fang, L., Zhang, Z., Ma, W., Niu, Y., Ju, L., Deng, J., Zhao, T., Lian, J., et al. (2019).
770 *Gossypium barbadense* and *Gossypium hirsutum* genomes provide insights into the origin and evolution
771 of allotetraploid cotton. *Nat. Genet.* *51*, 739-748.
- 772 Huang, J., Guo, Y., Sun, Q., Zeng, W., Li, J., Li, X., and Xu, W. (2019). Genome-wide identification of
773 R2R3-MYB transcription factors regulating secondary cell wall thickening in cotton fiber development.
774 *Plant Cell Physiol.* *60*, 687-701.

- 775 Ishida, T., Thitamadee, S., and Hashimoto, T. (2007). Twisted growth and organization of cortical
776 microtubules. *J. Plant Res.* *120*, 61-70.
- 777 Jaffe M. J., and Galston, A.W. (1968). The physiology of tendrils. *Annu. Rev. Plant Physiol.* *19*, 417-434.
- 778 Smyth, D.R. (2016). Helical growth in plant organs: mechanisms and significance. *Development* *143*,
779 3272-3282.
- 780 Jian, G., Ma, C., Zheng, C., (2003). Advances in cotton breeding for resistance to fusarium and
781 verticillium wilt in the last fifty years in China. *Agr. Sci. China* *2(3)*, 280-288
- 782 Joshi-Saha, A., and Reddy, K.S. (2015). Repeat length variation in the 5'UTR of myo-inositol
783 monophosphatase gene is related to phytic acid content and contributes to drought tolerance in chickpea
784 (*Cicer arietinum L.*). *J. Exp. Bot.* *66*, 5683-5690.
- 785 Kong, L., Cheng, J., Zhu, Y., Ding, Y., Meng, J., Chen, Z., Xie, Q., Guo, Y., Li, J., Yang, S., et al. (2015).
786 Degradation of the ABA co-receptor ABI1 by PUB12/13 U-box E3 ligases. *Nat. Commun.* *6*, 8630.
- 787 Kretzschmar, F.K., Mengel, L.A., Muller, A.O., Schmitt, K., Bliersch, K.F., Valerius, O., Braus, G.H., and
788 Ischebeck, T. (2018). PUX10 is a lipid droplet-localized scaffold protein that interacts with CELL
789 DIVISION CYCLE48 and is involved in the degradation of lipid droplet proteins. *Plant Cell* *30*, 2137-
790 2160.
- 791 Kumar, S., and Bhatia, S. (2016). A polymorphic (GA/CT)_n- SSR influences promoter activity of
792 tryptophan decarboxylase gene in *Catharanthus roseus L. Don.* *Sci. Rep.* *6*, 33280.
- 793 Lacape, J.M., Llewellyn, D., Jacobs, J., Arioli, T., Becker, D., Calhoun, S., Al-Ghazi, Y., Liu, S., Palai,
794 O., Georges, S., et al. (2010). Meta-analysis of cotton fiber quality QTLs across diverse environments in
795 a *Gossypium hirsutum* x *G. barbadense* RIL population. *BMC Plant Biol.* *10*, 132.
- 796 Lee, Y., Chou, T., Pittman, S.K., Keith, A.L., Razani, B., and Wehl, C.C. (2017). Keap1/Cullin3
797 modulates p62/SQSTM1 activity via UBA domain ubiquitination. *Cell Rep.* *19*, 188-202.
- 798 Li, F., Wu, S., Lü, F., Chen, T., Ju, M., Wang, H., Jiang, Y., Zhang, J., Guo, W., and Zhang, T. (2009).
799 Modified fiber qualities of the transgenic cotton expressing a silkworm fibroin gene. *Chinese Sci. Bull.*
800 *54*, 1210-1216.
- 801 Li, H., Sun, B., Sasabe, M., Deng, X., Machida, Y., Lin, H., Julie Lee, Y.R., and Liu, B. (2017).
802 *Arabidopsis* MAP65-4 plays a role in phragmoplast microtubule organization and marks the cortical cell
803 division site. *New Phytol.* *215*, 187-201.
- 804 Li, Y., Wang, N.N., Wang, Y., Liu, D., Gao, Y., Li, L., and Li, X.B. (2018). The cotton XLIM protein

- 805 (GhXLIM6) is required for fiber development *via* maintaining dynamic F-actin cytoskeleton and
806 modulating cellulose biosynthesis. *Plant J.* *96*, 1269-1282.
- 807 Liu, L., Zhang, Y., Tang, S., Zhao, Q., Zhang, Z., Zhang, H., Dong, L., Guo, H., and Xie, Q. (2010). An
808 efficient system to detect protein ubiquitination by agroinfiltration in *Nicotiana benthamiana*. *Plant J.* *61*,
809 893-903.
- 810 Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-time
811 quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* *25*, 402-408.
- 812 Lucas, J.R., and Shaw, S.L. (2012). MAP65-1 and MAP65-2 promote cell proliferation and axial growth
813 in *Arabidopsis* roots. *Plant J.* *71*, 454-463.
- 814 Ma, Q., Wang, N., Hao, P., Sun, H., Wang, C., Ma, L., Wang, H., Zhang, X., Wei, H., and Yu, S. (2019).
815 Genome-wide identification and characterization of TALE superfamily genes in cotton reveals their
816 functions in regulating secondary cell wall biosynthesis. *BMC Plant Biol.* *19*, 432.
- 817 Martinez-Sanz, M., Pettolino, F., Flanagan, B., Gidley, M.J., and Gilbert, E.P. (2017). Structure of
818 cellulose microfibrils in mature cotton fibres. *Carbohydr Polym.* *175*, 450-463.
- 819 McCouch SR, Chen X, Panaud O, Temnykh S, Xu Y, Cho YG, Huang N, Ishii T, Blair M. (1997).
820 Microsatellite marker development, mapping and applications in rice genetics and breeding. *Plant Mol.*
821 *Biol.* *35(1)*, 89-99.
- 822 Mei, M., Syed, N.H., Gao, W., Thaxton, P.M., Smith, C.W., Stelly, D.M., and Chen, Z.J. (2004). Genetic
823 mapping and QTL analysis of fiber-related traits in cotton (*Gossypium*). *Theor. Appl. Genet.* *108*, 280-
824 291.
- 825 Mirkin, S.M. (2007). Expandable DNA repeats and human disease. *Nature* *447*, 932-940.
- 826 Nakajima, K., Furutani, I., Tachimoto, H., Matsubara, H., and Hashimoto, T. (2004). SPIRAL1 encodes
827 a plant-specific microtubule-localized protein required for directional control of rapidly expanding
828 *Arabidopsis* cells. *Plant Cell* *16*, 1178-1190.
- 829 Neuber, O., Jarosch, E., Volkwein, C., Walter, J., and Sommer, T. (2005). Ubx2 links the Cdc48 complex
830 to ER-associated protein degradation. *Nat. Cell Biol.* *7*, 993-998.
- 831 Ning, Z., Chen, H., Mei, H., and Zhang, T. (2014). Molecular tagging of QTLs for fiber quality and yield
832 in the upland cotton cultivar Acala-Prema. *Euphytica* *195*, 143-156.
- 833 Ooijen, J., and Voorrips, R. E. (2001). JoinMap version 3.0: software for the calculation of genetic linkage
834 maps. *Int. J. Plant Res.*

- 835 Paredez, A.R., Somerville, C.R., and Ehrhardt, D.W. (2006). Visualization of cellulose synthase
836 demonstrates functional association with microtubules. *Science* 312, 1491-1495.
- 837 Paterson, A.H., Brubaker, C.L., and Wendel, J.F. (1993). A rapid method for extraction of cotton
838 (*Gossypium spp.*) genomic DNA suitable for RFLP or PCR analysis. *Plant Mol. Biol. Rep.* 11, 122-127.
- 839 Qin, Z., Bai, Y., Muhammad, S., Wu, X., Deng, P., Wu, J., An, H., and Wu, L. (2019). Divergent roles of
840 FT-like 9 in flowering transition under different day lengths in *Brachypodium distachyon*. *Nat. Commun.*
841 10, 812.
- 842 Ranathunge, C., Wheeler, G.L., Chimahusky, M.E., Kennedy, M.M., Morrison, J.I., Baldwin, B.S.,
843 Perkins, A.D., and Welch, M.E. (2018). Transcriptome profiles of sunflower reveal the potential role of
844 microsatellites in gene expression divergence. *Mol. Ecol.* 27, 1188-1199.
- 845 Rong, J., Feltus, F.A., Waghmare, V.N., Pierce, G.J., Chee, P.W., Draye, X., Saranga, Y., Wright, R.J.,
846 Wilkins, T.A., May, O.L., et al. (2007). Meta-analysis of polyploid cotton QTL shows unequal
847 contributions of subgenomes to a complex network of genes and gene clusters implicated in lint fiber
848 development. *Genetics* 176, 2577-2588.
- 849 Sambade A, Findlay K, Schäffner AR, Lloyd CW, Buschmann H. (2014). Actin-dependent and -
850 independent functions of cortical microtubules in the differentiation of *Arabidopsis* leaf trichomes. *Plant*
851 *Cell* 26, 1629-1644.
- 852 Sedbrook, J.C., Ehrhardt, D.W., Fisher, S.E., Scheible, W.R., and Somerville, C.R. (2004). The
853 *Arabidopsis* sku6/spiral1 gene encodes a plus end-localized microtubule-interacting protein involved in
854 directional cell expansion. *Plant Cell* 16, 1506-1520.
- 855 Schuberth, C., and Buchberger, A. (2005). Membrane-bound Ubx2 recruits Cdc48 to ubiquitin ligases
856 and their substrates to ensure efficient ER-associated protein degradation. *Nat. Cell Biol.* 7, 999-1006.
- 857 Shoji, T., Narita, N.N., Hayashi, K., Asada, J., Hamada, T., Sonobe, S., Nakajima, K., and Hashimoto, T.
858 (2004). Plant-specific microtubule-associated protein SPIRAL2 is required for anisotropic growth in
859 *Arabidopsis*. *Plant Physiol.* 136, 3933-3944.
- 860 Skotheim, J.M., and Mahadevan, L. (2005). Physical limits and design principles for plant and fungal
861 movements. *Science* 308, 1308-1310.
- 862 Sun, Q., Huang, J., Guo, Y., Yang, M., Guo, Y., Li, J., Zhang, J., and Xu, W. (2020). A cotton NAC
863 domain transcription factor, GhFSN5, negatively regulates secondary cell wall biosynthesis and anther
864 development in transgenic *Arabidopsis*. *Plant Physiol. Bioch.* 146, 303-314.

- 865 Tiwari, S. C. , Wilkins, T. A. . (1995). Cotton (*Gossypium hirsutum*) seed trichomes expand via diffuse
866 growing mechanism. *Can. J. Bot.* *73*, 746-775
- 867 Ulloa, M., and M. Wrjr. (2000). Genetic linkage map and QTL analysis of agronomic and fiber quality
868 traits in an intraspecific population. *J. Cotton Sci.* *4(3)*, 161-170.
- 869 Varshavsky A. (2017). The Ubiquitin System, Autophagy, and Regulated Protein Degradation. *Annu. Rev.*
870 *Biochem.* *86*, 123-128.
- 871 Voorrips, R.E. (2002). MapChart 2.2: software for the graphical presentation of linkage maps and QTLs.
872 *J. Hered.* *93(1)*, 77-78.
- 873 Wang, S., Kurepa, J., Hashimoto, T., and Smalle, J.A. (2011). Salt stress-induced disassembly of
874 *Arabidopsis* cortical microtubule arrays involves 26S proteasome-dependent degradation of SPIRAL1.
875 *Plant Cell* *23*, 3412-3427.
- 876 Wang, Y., Ning, Z., Hu, Y., Chen, J., Zhao, R., Chen, H., Ai, N., Guo, W., and Zhang, T. (2015). Molecular
877 mapping of restriction-site associated DNA markers in allotetraploid upland cotton. *PLoS ONE* *10*,
878 e0124781.
- 879 Wu, S., Wang, H., Li, F., Chen, T., Zhang, J., Jiang, Y., Ding, Y., Guo, W., and Zhang, T. (2008). Enhanced
880 agrobacterium-mediated transformation of embryogenic calli of upland cotton via efficient selection and
881 timely subculture of somatic embryos. *Plant Mol. Biol. Rep.* *26*, 174-185.
- 882 Xia, T., Li, N., Dumenil, J., Li, J., Kamenski, A., Bevan, M.W., Gao, F., and Li, Y. (2013). The ubiquitin
883 receptor DA1 interacts with the E3 ubiquitin ligase DA2 to regulate seed and organ size in *Arabidopsis*.
884 *Plant Cell* *25*, 3347-3359.
- 885 Yanagisawa M, Desyatova AS, Belteton SA, Mallery EL, Turner JA, Szymanski DB. (2015). Patterning
886 mechanisms of cytoskeletal and cell wall systems during leaf trichome morphogenesis. *Nature Plants* *1*,
887 15014.
- 888 Yang, Y., Fu, D., Zhu, C., He, Y., Zhang, H., Liu, T., Li, X., and Wu, C. (2015). The RING-Finger
889 ubiquitin ligase HAF1 mediates heading date 1 degradation during photoperiodic flowering in rice. *Plant*
890 *Cell* *27*, 2455-2468.
- 891 Zhang, J., Huang, G.Q., Zou, D., Yan, J.Q., Li, Y., Hu, S., and Li, X.B. (2018). The cotton (*Gossypium*
892 *hirsutum*) NAC transcription factor (FSN1) as a positive regulator participates in controlling secondary
893 cell wall biosynthesis and modification of fibers. *New Phytol.* *217*, 625-640.
- 894 Zhang, M., Yu, Q., Liu, Z., Liang, C., Zhang, B., and Li, M. (2017). UBX domain-containing proteins

895 are involved in lipid homeostasis and stress responses in *Pichia pastoris*. *Int. J. Biochem. Cell Biol.* *90*,
896 136-144.

897 Zhang, T., Yuan, Y., Yu, J., Guo, W., and Kohel, R.J. (2003). Molecular tagging of a major QTL for fiber
898 strength in upland cotton and its marker-assisted selection. *Theor. Appl. Genet.* *106*, 262-268.

899

900

Journal Pre-proof

Highlights

- Isolation of the first fiber strength gene GhUBX using map-based cloning strategy.
- Verification of the function of GhUBX experimentally in transgenic cotton.
- Link helices to the cotton fiber strength, that more helices make fiber stronger.
- An ubiquitin–proteasome system regulating the development of cotton fiber.

